

Thermal rectification in quantum graded mass systems

Emmanuel Pereira

Departamento de Física-ICEx, UFMG, CP 702, 30.161-970 Belo Horizonte MG, Brazil

e-mail: emmanuel@fisica.ufmg

March 1, 2010

Abstract

We show the existence of thermal rectification in the graded mass quantum chain of harmonic oscillators with self-consistent reservoirs. Our analytical study allows us to identify the ingredients leading to the effect. The presence of rectification in this effective, simple model (representing graded mass materials, systems that may be constructed in practice) indicates that rectification in graded mass quantum systems may be an ubiquitous phenomenon. Moreover, as the classical version of this model does not present rectification, our results show that, here, rectification is a direct result of the quantum statistics.

PACS number(s): 05.70.Ln; 05.40.-a; 44.10.+i

Key Words: heat flow; thermal rectification; quantum graded mass systems

A fundamental challenge in statistical physics is the derivation of macroscopic phenomenological laws of thermodynamic transport from the underlying microscopic Hamiltonian systems. However, after decades, a first-principle derivation of the Fourier's law of heat conduction, for instance, is still a puzzle [1]. Many works have been devoted to the theme [2], most of them by means of computer simulations. But, in these problems where the central question involves the convergence or divergence of thermal conductivity, sometimes there is a significant difficulty to arrive at precise conclusions from numerical results [3]. Thus, the necessity of analytical studies together with the huge complexity of the associated nonlinear dynamical systems led to several works considering approximative schemes or simplified models [4] - since the pioneering work of Debye, the microscopic models to describe heat conduction are mainly given by systems of anharmonic oscillators, leading to problems without analytical solutions. Many works involve, e.g., the use of approximations such as Boltzmann equations, master equations for effective models, the analysis of Green-Kubo formula, etc. An example of simple (effective) model that can be analytically studied is the classical or quantum harmonic chain of oscillators with self-consistent reservoirs. It has been proposed a while ago [5, 6], but it is always revisited [7, 8]. In such model, each site of the chain is coupled to a reservoir; the first and last sites are coupled to "real" thermal baths, while the inner reservoirs only mimic the absent anharmonic interactions. The self-consistent condition means that there is no heat flow between an inner reservoir and its site in the steady state: the inner baths act only as a phonon scattering mechanism, such as

the on-site anharmonic potential in more elaborated systems. This model, the classical and also the quantum version, in opposition to a standard harmonic system, obeys the Fourier's law [7].

In this scenario of intensive study of the heat mechanism, problems involving the possibility to control the heat flow by constructing thermal nano-instruments such as thermal diodes, and even transistors, thermal gates and memories, have recently attracted considerable theoretical [9, 10, 11] and experimental [12] interest. A thermal diode, or rectifier, is a device in which the magnitude of the heat current changes as we invert the device between two thermal baths. That is, in a thermal diode, heat flows preferably in one direction. There are some analytical attempts to explain the phenomenon of thermal rectification and/or design a diode by using simple models (see e.g. the spin-boson nanojunction [13], [14], and the billiard system [15]) but, again, most of the works consider numerical computations [9, 10, 11]. It is worth to recall the extensive work of B. Li and collaborators: e.g. thermal rectification in asymmetric graphene ribbons is studied by using molecular dynamics simulations in ref. [16]; also in carbon nanocone structures [17]; at silicon-amorphous polyethylene interfaces [18]; in carbon nanotube intramolecular junctions [19], etc. A commonly used design of diodes is given by the sequential coupling of two or three chains with different anharmonic potentials [9, 10, 11]. It is frequently studied, but it is also criticized due to the difficulty to be constructed in practice [10]. Recently, Chang et al. [12], considering a different procedure, built a nanoscale thermal rectifier in an experimental work: they use a graded material, namely, nanotubes externally and inhomogeneously mass-loaded with heavy molecules, a system with asymmetric-axial thermal conductance. Graded materials are also considered in other numerical studies of diodes [20, 21], and the interesting results made some authors suggest that thermal rectification is guaranteed by the use of graded materials with anharmonic interactions.

Having in mind the investigation of this suggested relation between graded (anharmonic) materials and rectification, in a recent paper [22] we study the classic harmonic chain of oscillators with self-consistent reservoirs (CSC). We show the absence of thermal rectification for the CSC model with graded masses or graded interparticle interactions by using the method developed in refs.[23], [24]. In [25], Segal obtains similar results by using the approach of [8]. We, and other authors [5, 7], understand the SC models as effective anharmonic models (the inner reservoirs mimic the absent anharmonic potentials), but it is not a consensus. Anyway, even if we discard any relation with anharmonicity, these results show, at least, that the onset of Fourier's law does not guarantee thermal rectification in asymmetric classic models - we recall that the Fourier's law does not hold in standard harmonic systems (i.e., harmonic models with reservoirs at the boundaries only), but it holds in the SC models.

Still searching for the mechanism behind the rectification in graded mass systems, we recall that, at low temperatures, quantum effects may introduce significant changes. The thermal conductivity for the quantum self-consistent harmonic chain (QSC), for example, depends on temperature [8, 26], in opposition to the CSC thermal conductivity [7]. Thus, searching for possible quantum effects in the mechanism of thermal rectification, in a quite recent work [27], we turn to the inhomogeneous QCS and, in the linear response regime, i.e., for the chain submitted to a very small gradient of temperature and considering only linear corrections in computations with changes in the temperature, we show the absence of thermal rectification despite quantum effects in the conductivity. However, we may still ask about the possibility of a rectification for the quantum system submitted to a large gradient of temperature or, at least, for some effect beyond linear corrections. It is, indeed, a very important question: in ref.[28], the authors show that there is a significative rectification in some chaotic billiard systems "provided the temperatures (of the two sides of the system) are strongly different..."

In the present work, with the focus on the onset of rectification in inhomogeneous systems, we revisit the graded mass QSC model, and study it by using an analytical approach valid for any temperature gradient, i.e., beyond the linear response regime. For this quantum model, we show that, in opposition to the behavior of its classical version, there is a thermal rectification. We offer an explanation for this difference: in the quantum version, the expression for the heat flow (3) involves a distribution for the phonon frequencies (the frequencies in the system) that depends on temperature (5), leading to a mix of temperature, frequencies and also particle masses (see eqs.(3, 4)). It does not happen in the classical system - see the comments for the high temperature limit at the end of this manuscript. This intricate mix of temperature, masses and frequencies makes the heat current change as we invert the graded mass system between two thermal reservoirs. The presence of rectification in this quite simple, say, bare model is of considerable interest: it indicates the generality of such phenomenon in the experimentally realizable class of graded mass systems, i.e., it indicates that rectification is not related to intricate interactions in the microscopic systems. Moreover, as far as we know, this work is the first demonstration of a “purely” quantum thermal rectifier, in the sense that, for such model, the rectification is a direct result of the quantum statistics: i.e., rectification is absent in the classical version of the model.

Let us introduce the QSC model. We use a Ford-Kac-Mazur approach, as detailed presented in ref.[8], in order to describe the quantum system and its time evolution to the steady state. Here, all the baths connected to the chain are modeled as mechanical harmonic systems, with initial coordinates and momenta determined by some statistical distribution. Then, we solve the quantum dynamics given by Heisenberg equations, take the stochastic distribution for the initial coordinates of the baths, as well as the limit $t \rightarrow \infty$, and obtain the expression for the heat flow in the steady state.

The Hamiltonian of our system, a chain (W) with harmonic interparticle and on-site potentials, with each site connected to a bath (B), also with harmonic interactions, is given by

$$\begin{aligned}\mathcal{H} &= \mathcal{H}_W + \sum_{i=1}^N \mathcal{H}_{B_i} + \sum_{i=1}^N X_W^T V_{B_i} X_{B_i}, \\ \mathcal{H}_S &= \frac{1}{2} \dot{X}_S^T M_S \dot{X}_S + \frac{1}{2} X_S^T \Phi_S X_S,\end{aligned}\tag{1}$$

where $S = W$ or B_i ; M_W , M_{B_i} are the particle-mass diagonal matrices for the chain and baths; Φ_W and Φ_{B_i} are symmetric matrices describing the interparticle and on-site harmonic interactions; and V_{B_i} gives the interaction between the i -site and its bath (more details below). We have, for each part, $X = [X_1, X_2, \dots, X_{N_s}]^T$, where X_r is the position operator of the r -th particle; $\dot{X} = M^{-1}P$, where P_l is the momentum operator of the l -th particle. Of course, it follows that $[X_r, P_l] = i\hbar\delta_{r,l}$. Finally, the dynamics is given by the Heisenberg equations

$$\begin{aligned}M_W \ddot{X}_W &= -\Phi_W X_W - \sum_i V_{B_i}^T X_{B_i}, \\ M_{B_i} \ddot{X}_{B_i} &= -\Phi_{B_i} X_{B_i} - \sum_i V_{B_i}^T X_W.\end{aligned}\tag{2}$$

The formulas for the heat currents inside the chain and from each reservoir to the chain are related to $\langle X_W \dot{X}_W^T \rangle$ and $\langle X_B \dot{X}_W^T \rangle$. Before presenting the formulas, let us give a very short resume of their derivation. To find the expressions, we turn to the Heisenberg equations (2), treat the equations of the baths as linear inhomogeneous equations, and plug these solutions

into the equations for the chain. Then, we take the average over the initial conditions of the baths, which are assumed to be distributed according to equilibrium phonon distributions with properly chosen temperatures, determined such that the self-consistent condition holds, that is, we must take the temperatures of the inner baths such that there is no heat flow between an inner bath and its site. We reach the steady state by taking the limit $t \rightarrow \infty$. For technical reasons we still take $t_0 \rightarrow -\infty$, and consider the Fourier transform of t . We note that, by plugging the solutions of the baths back into the equations of motion for the system, we get a quantum Langevin equation. Remember that, for our model, all particles are connected to heat reservoirs that are taken to be Ohmic. The coupling strength to the reservoirs is controlled by the dissipation constant ζ defined by an expression involving the matrix V presented in the Hamiltonian above (see ref.[8]).

The expression for the heat flow from the l -th reservoir to the chain is given by

$$\begin{aligned} \mathcal{F}_l &= \sum_{m=1}^N \zeta^2 \int_{-\infty}^{+\infty} d\omega \, \omega^2 \left| [G_W(\omega)]_{l,m} \right|^2 \times \\ &\times \frac{\hbar\omega}{\pi} [f(\omega, T_l) - f(\omega, T_m)], \end{aligned} \quad (3)$$

where ζ is the dissipation constant;

$$[G_W(\omega)]^{-1} = -\omega^2 M_W + \Phi_W - \sum_l \Sigma_l^+(\omega), \quad (4)$$

the matrix Σ_l^+ above has only one non-vanishing element: $(\Sigma_l^+)_{l,l} = i\zeta\omega$; $f(\omega, T_l)$ is the phonon distribution for the l -th bath

$$f(\omega, T_l) = 1/[\exp(\hbar\omega/KT_l) - 1]; \quad (5)$$

the variable ω in the expressions above comes from the Fourier transform, namely, $\tilde{X}(\omega) = (2\pi)^{-1} \int_{-\infty}^{+\infty} dt \, X(t) \, e^{i\omega t}$. We stress that a detailed derivation of the heat flow formula (3) is presented in ref.[8].

The heat flow in the chain may be given by $\mathcal{F}_1$, i.e., by the heat flow from the first reservoir, together with the self-consistent condition $\mathcal{F}_2 = \mathcal{F}_3 = \dots = \mathcal{F}_{N-1} = 0$, which means that the inner reservoirs do not inject energy into the system (the energy comes from the first reservoir, passes through the chain and goes out to the last reservoir). In such condition, we have $\mathcal{F}_1 = -\mathcal{F}_N$. To investigate the presence or absence of thermal rectification, we need to examine the heat current in a chain with reservoirs at temperatures $T_1, T_2, \dots, T_N$, where T_1 and T_N are given, and $T_2, \dots, T_{N-1}$ are determined by the self-consistent condition ($\mathcal{F}_2 = 0, \dots, \mathcal{F}_{N-1} = 0$); and also in the same chain with inverted thermal reservoirs, precisely, with the baths taken at temperatures $T'_1, T'_2, \dots, T'_N$, where $T'_1 = T_N$, $T'_N = T_1$, and $T'_2, \dots, T'_{N-1}$ are determined by the self-consistent condition ($\mathcal{F}'_2 = 0, \dots, \mathcal{F}'_{N-1} = 0$). In the case of absence of thermal rectification, we shall obtain $\mathcal{F}_1 + \mathcal{F}'_1 = 0$. That is what we will examine.

The determination of $G_W(w)$, the matrix given by eq. (4) and necessary to study $\mathcal{F}_l$, may be a very difficult task – we know a precise solution for some specific cases: e.g., for a homogeneous next-neighbors interparticle interaction and all particles with the same mass $M_W = mI$ [8]; and also for the case of particles with alternate masses, i.e. $m_j = m_1$ for j odd, and $m_j = m_2$ for j even [26]. Hence, we first restrict our interparticle interaction Φ to the homogeneous nearest-neighbor case, i.e., $\Phi = -\Delta$, the lattice Laplacian, and study the equations for the smallest

possible chain: $N = 3$ (it is, in fact, a cell of a large chain). Then, we show that a graded mass distribution leads to a thermal rectification. After that, we argue to show that similar relations extend to $N = 4$, and so on.

Thus, for $N = 3$, we have

$$G_W^{-1}(w) = \begin{bmatrix} R_{1,1} & -1 & 0 \\ -1 & R_{2,2} & -1 \\ 0 & -1 & R_{3,3} \end{bmatrix}, \quad (6)$$

where $R_{k,k} \equiv g_{k,k}^{-1} = 2 - w^2 M_k - i\zeta w$. To have graded masses, we take $M_1 < M_2 < M_3$. And using the notation $c \equiv \zeta^2 \hbar / \pi$, $A_{lm} \equiv |(G_W(w))_{l,m}|^2$, $f_k \equiv f(w, T_k)$, we get (from eq.(3))

$$\begin{aligned} \mathcal{F}_1 &= c \int dw w^3 A_{12} [f_1 - f_2] + c \int dw w^3 A_{13} [f_1 - f_3], \\ \mathcal{F}_3 &= c \int dw w^3 A_{31} [f_3 - f_1] + c \int dw w^3 A_{32} [f_3 - f_2]. \end{aligned}$$

It is easy to see that $A_{jk} = A_{kj}$. By taking the inverse of G_W^{-1} above, we have

$$\begin{aligned} A_{12} &= [(2 - w^2 M_3)^2 + \zeta^2 w^2] / |\det G_W^{-1}|^2, \\ A_{32} &= [(2 - w^2 M_1)^2 + \zeta^2 w^2] / |\det G_W^{-1}|^2. \end{aligned} \quad (7)$$

Note that, by using $\mathcal{F}_1 = -\mathcal{F}_3$, from the equations above we obtain (as expected) $\mathcal{F}_2 = 0$. For the chain with inverted reservoirs, we obtain the expressions for $\mathcal{F}'_1$ and $\mathcal{F}'_3$ similar to those for $\mathcal{F}_1$ and $\mathcal{F}_3$ above, but with the change $T_1 \leftrightarrow T_3$, i.e., with the replacement $f_1 \leftrightarrow f_3$, and $f_2 \rightarrow f'_2$. The question is: do we have $\mathcal{F}_1 + \mathcal{F}'_1 = 0$?

From $\mathcal{F}_1 = -\mathcal{F}_3$ (i.e., $\mathcal{F}_2 = 0$) and $\mathcal{F}'_1 = -\mathcal{F}'_3$ (i.e., $\mathcal{F}'_2 = 0$), we obtain

$$\int dw w^3 A_{12} [f_1 - f_2] = \int dw w^3 A_{32} [f_2 - f_3], \quad (8)$$

$$\int dw w^3 A_{12} [f_3 - f'_2] = \int dw w^3 A_{32} [f'_2 - f_1]. \quad (9)$$

To get the absence of rectification, we must have $\mathcal{F}_1 + \mathcal{F}'_1 = 0$, i.e.,

$$\int dw w^3 A_{12} [f_1 - f_2] = \int dw w^3 A_{12} [f'_2 - f_3], \quad (10)$$

or, which comes directly from eqs.(8,9,10) (and also from $\mathcal{F}_3 + \mathcal{F}'_3 = 0$)

$$\int dw w^3 A_{32} [f_3 - f_2] = \int dw w^3 A_{32} [f'_2 - f_1]. \quad (11)$$

That is, if we have eqs.(8,9,10), then eq.(11) follows; similarly, given eqs.(8,9,11), we have eq.(10).

Let us use the short notation

$$F_1(T_k) = \int dw w^3 A_{12} f_k, \quad F_2(T_k) = \int dw w^3 A_{32} f_k,$$

and similarly for $F_j(T'_2)$. From eqs.(8) and (9) we obtain

$$F_1(T_2) + F_2(T_2) = F_1(T_1) + F_2(T_3), \quad (12)$$

$$F_1(T'_2) + F_2(T'_2) = F_2(T_1) + F_1(T_3). \quad (13)$$

We write the condition for absence of rectification (10) as

$$F_1(T_2) + F_1(T'_2) = F_1(T_1) + F_1(T_3), \quad (14)$$

or, which comes from these equations,

$$F_2(T_2) + F_2(T'_2) = F_2(T_1) + F_2(T_3). \quad (15)$$

Let us briefly give an idea of our analysis before turning to the mathematical expressions. It follows that any function $F_j(T)$ above, and also the sum $F_1(T) + F_2(T)$, is a monotone, strictly increasing, convex function of T (it is easily verified by studying its derivative d/dT). Hence, given T_1 and T_3 , the value of T_2 is univocally determined by eq.(12), and the value of T'_2 by eq.(13). Moreover, $F_1(T)$ is bigger and increases faster than $F_2(T)$, for $M_1 < M_2 < M_3$. In short, we have three different equations for T_2 and T'_2 , that are determined by the two first equations; as the third one may not be derived (it is independent) from the first and second equations, we shall have rectification, i.e. the third equation will not be always satisfied. For linear equations in T (that is the case for the approximations of high temperature or linear response regime, as we show ahead), the system of two variables and three equations has a solution only if it is linearly dependent - more details ahead.

Now we make explicit, i.e., we give the mathematical details of the analysis described above for the eqs.(12, 13, 14). For simplicity, we will consider small temperatures (and so, also small changes in T), but, of course, we will go beyond linear corrections. Since F (F_1 and F_2) is an analytic function of T (for $T > 0$), we write, for some $T_0 \neq 0$, $|T_0 - T|$ small,

$$\begin{aligned} F_1(T) &= F_1(T_0) + F'_1(T_0)(T - T_0) + \frac{F''_1(T_0)}{2}(T - T_0)^2 \\ &= \left[F_1(T_0) - F'_1(T_0)T_0 + \frac{F''_1(T_0)}{2}T_0^2 \right] + \left[F'_1(T_0) - 2T_0 \frac{F''_1(T_0)}{2} \right] T + \frac{F''_1(T_0)}{2}T^2 \\ &\equiv a_0 + a_1T + a_2T^2, \end{aligned} \quad (16)$$

that is, we discard corrections up to $\mathcal{O}((T - T_0)^3)$. We have (at least for small T_0) $a_1, a_2 > 0$. The same follows for $F_2(T)$,

$$F_2(T) = b_0 + b_1T + b_2T^2. \quad (17)$$

Hence, introducing eqs.(16, 17) in eq.(12), we get, for small T_1 and T_3 ,

$$T_2 = \frac{-(a_1 + b_1) \pm \sqrt{(a_1 + b_1)^2 + 4(A_1 + B_3)(a_2 + b_2)}}{2(a_2 + b_2)}, \quad (18)$$

where

$$A_1 \equiv a_1T_1 + a_2T_1^2, \quad B_3 \equiv b_1T_3 + b_2T_3^2. \quad (19)$$

Of course, only $T_2 > 0$ makes sense, and so, T_2 is determined by the positive expression in eq.(18) above. Similarly, for T'_2 we get, from eq.(13),

$$T'_2 = \frac{-(a_1 + b_1) \pm \sqrt{(a_1 + b_1)^2 + 4(A_3 + B_1)(a_2 + b_2)}}{2(a_2 + b_2)}, \quad (20)$$

where

$$A_3 \equiv a_1T_3 + a_2T_3^2, \quad B_1 \equiv b_1T_1 + b_2T_1^2. \quad (21)$$

Again, we consider the positive expression for T'_2 .

Now we check if eq.(14) is verified or not, that means absence or existence of rectification respectively. Namely, we check if it is true the equality

$$(a_1 T_2 + a_2 T_2^2) + (a_1 T'_2 + a_2 T'^2_2) = (a_1 T_1 + a_2 T_1^2) + (a_1 T_3 + a_2 T_3^2). \quad (22)$$

To make the investigation, we introduce the (positive) expressions for T_2 (18) and T'_2 (20) in equation above. By taking $T_1 = T$, $T_3 = T + \epsilon$, with small ϵ , and using

$$\sqrt{1+x} = 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \mathcal{O}(x^3),$$

we easily verify that eq.(22) is not satisfied. E.g., by taking in eq.(22) the terms involving only ϵ and ϵ^2 (not those with T , $T\epsilon$, etc), we get, for the L.H.S. and R.H.S. of eq.(22),

$$\begin{aligned} \text{L.H.S.} &= a_1 \epsilon + \frac{a_1}{a_1 + b_1} (a_2 + b_2) \epsilon^2 + (a_2 b_1 - a_1 b_2) \frac{(b_1^2 + a_1^2)}{(a_1 + b_1)^3} \epsilon^2, \\ \text{R.H.S.} &= a_1 \epsilon + a_2 \epsilon^2. \end{aligned}$$

Note that there is an equality up to first order in ϵ , and so, as we have already recalled [27], we do not see rectification in the linear regime. But, considering the corrections beyond the linear approximation, the difference is evident, which shows the existence of rectification.

For transparency, let us also give more details of our analysis for some situations of absence of rectification. It happens, e.g., in the following cases. (I) For the system with homogeneous mass distribution (obviously): if $M_1 = M_2 = M_3$, then $A_{12} = A_{32}$ (and so, eq.(14) is obtained by the sum of eqs.(12) and (13)). (II) In the high temperature regime: in such case, as shown in ref.[8], the quantum system behaves as the classical one, where there is no rectification (as already shown in refs.[22] and [25]). To see this result from our previous equations, note that for large T we have $w/[\exp(\hbar w/KT) - 1] \rightarrow KT/\hbar$, and so, eqs.(12), (13), and (14) become linear equations in T_2 , T'_2 , T_1 and T_3 . It is easy to see that they are linearly dependent (and so, eq.(14) comes from eqs.(12) and (13)). (III) In the linear response regime (already shown in ref.[27] and commented above): again, as described in ref.[8], for small temperature gradients we have

$$\mathcal{F}_k = c \int dw w^3 \frac{\partial f}{\partial T}(w, T) \sum_{m=1}^3 A_{km} (T_k - T_m),$$

where T above is some average temperature. The expression above leads us to the same case of (II) (i.e., we have linear equations in T_2 , T'_2 , etc).

Now we turn to larger chains. For $N = 4$, we have the equations $\mathcal{F}_1 = -\mathcal{F}_4$ and $\mathcal{F}_2 = 0$, $\mathcal{F}_3 = 0$ (the self-consistent equation). But, as $\mathcal{F}_1 + \mathcal{F}_2 + \mathcal{F}_3 + \mathcal{F}_4 = 0$ (it may be easily verified from the expressions for $\mathcal{F}$), if we take $\mathcal{F}_1 = -\mathcal{F}_4$ and $\mathcal{F}_2 = 0$, then $\mathcal{F}_3 = 0$ follows (i.e., it is not a new equation; and vice-versa: $\mathcal{F}_2 = 0$ and $\mathcal{F}_3 = 0$ leads to $\mathcal{F}_1 = -\mathcal{F}_4$). Given T_1 and T_4 , these two equations (e.g., $\mathcal{F}_1 = -\mathcal{F}_4$ and $\mathcal{F}_2 = 0$) determines T_2 and T_3 . The same follows for the system with inverted reservoirs (T'_2 and T'_3). Hence, the equation for the absence of rectification $\mathcal{F}_1 + \mathcal{F}'_1 = 0$ is an extra equation involving variables already determined, and the analysis follows as before (for $N = 4$ and any other N). We emphasize that the study of the dependence of the rectification on the system size (not presented here) is of importance: recall that, in the two-segment diode, the rectification vanishes as the system size goes to infinite.

We note that, for a system with three sites, if we take a mass distribution with $M_1 \neq M_3$ (e.g., $M_1 = M_2 \neq M_3$, or $M_1 \neq M_2 = M_3$) we still have rectification. For larger N , the analysis for the graded system is “automatic”, but it may be intricate for any other mass distribution. It is an interesting (and difficult) problem to investigate which inhomogeneity in the mass distribution is enough to guarantee the rectification.

We still recall that, for a standard harmonic system, i.e., for the harmonic chain with reservoirs at the boundaries only (in other words, turning off the inner reservoirs, say, the “schematic anharmonicity”), we have (see ref.[8]),

$$\mathcal{F}_1 = c \int dw w^3 A_{1N}(f_1 - f_N),$$

and similarly for $\mathcal{F}'$, with the replacement $f_1 \leftrightarrow f_N$; i.e., $\mathcal{F}_1 = -\mathcal{F}'_1$, and so, there is no rectification.

To conclude, our results show the existence of thermal rectification in the graded mass QSC, a quite simple model. It indicates that the presence of some phonon scattering mechanism, or some effective anharmonicity, is enough to guarantee the thermal rectification in an inhomogeneous quantum model. We stress the importance of the quantum nature of the reservoirs that provides an energy mixing (5). It is probable that some other intricate mechanism, such as real anharmonic potentials, may provide such mix in classical models, as indicated by numerical studies [20]. Finally, we recall that graded mass models are not only theoretical systems: they may be constructed in practice [12]. In short, diodes of graded mass systems sound to be ubiquitous (and experimentally reliable) structures.

Acknowledgments

We are in debt to the referee of a previous (unpublished) work to point out a mistake in our tentative to show the absence of thermal rectification for the QSC model. We also thank the reviewers of the present manuscript for several comments that improved its presentation. Work supported by CNPq and Fapemig (Brazil).

References

- [1] F. Bonetto, J. L. Lebowitz, L. Rey-Bellet, Mathematical Physics 2000, Imperial College Press (2000) pp. 128-150.
- [2] S. Lepri, R. Livi, A. Politi, Phys. Rep. 377 (2003) 1-80.
- [3] O. V. Gendelman, A. V. Savin, Phys. Rev. Lett. 92 (2004) 074301.
- [4] A. Dhar, Adv. Phys. 57 (2008) 457.
- [5] M. Bosterli, M. Rich, W.M. Visscher, Phys. Rev. A 1 (1970) 1086.
- [6] W.M. Visscher, M. Rich, Phys. Rev. A 12 (1975) 675.
- [7] F. Bonetto, J.L. Lebowitz, J. Lukkarinen, J. Stat.Phys 116 (2004) 783.
- [8] A. Dhar, D. Roy, J. Stat. Phys. 125 (2006) 801.

- [9] M. Terraneo, M. Peyrard, G. Casati, Phys. Rev. Lett. 88, (2002) 094302.
- [10] B. Hu, L. Yang, Y. Zhang, Phys. Rev. Lett. 97 (2006) 124302.
- [11] B. Li, L. Wang, G. Casati, Phys. Rev. Lett. 93 (2004) 184301.
- [12] C.W. Chang, D. Okawa, A. Majumdar, A. Zettl, Science 314 (2006) 1121.
- [13] D. Segal, A. Nitzan, Phys. Rev. Lett. 94 (2005) 034301.
- [14] L.-A. Wu, D. Segal, Phys. Rev. Lett. 102 (2009) 095503.
- [15] J.P. Eckmann, C. Mejía-Monasterio, Phys. Rev. Lett. 97 (2006) 094301.
- [16] N. Yang, G. Zhang, B. Li, Appl. Phys. Lett. 95 (2009) 033107.
- [17] N. Yang, G. Zhang, B. Li, Appl. Phys. Lett. 93 (2008) 243111.
- [18] M. Hu, P. Keblinski, B. Li, Appl. Phys. Lett. 92 (2008) 211908.
- [19] G. Wu, B. Li, Phys. Rev. B 76 (2007) 085424.
- [20] N. Yang, N. Li, L. Wang, B. Li, Phys. Rev. B 76 (2007) 020301 (R).
- [21] G. Casati, Nat. Nanotechnol. 2 (2007) 23.
- [22] E. Pereira, H. C. F. Lemos, Phys. Rev. E 78 (2008) 031108.
- [23] E. Pereira, R. Falcao, Phys. Rev. E 70 (2004) 046105.
- [24] E. Pereira, R. Falcao, Phys. Rev. Lett. 96 (2006) 100601.
- [25] D. Segal, Phys. Rev. E 79, (2009) 012103.
- [26] A. Francisco Neto, H.C.F. Lemos, E. Pereira, Phys. Rev. E 76 (2007) 031116.
- [27] E. Pereira, H. C. F. Lemos, J. Phys. A: Math. Theor. 42 (2009) 225006.
- [28] G. Casati, C. Mejía-Monasterio, T. Prosen, Phys. Rev. Lett. 98 (2007) 104302.